Total number of printed pages-7 villabl sies nego owi lo noi 3 (Sem-3/CBCS) STA HC 3 is open. (ii) Every open set is an union of open STATISTICS (Honours Core) Paper: STA-HC-3036 (Mathematical Analysis) The set of all integers is countable. Full Marks: 60 State Bolzano W. erstrass theorem. The figures in the margin indicate full marks for the questions. Answer the following questions as directed: $7 = 7 \times 1$ The value of $\Lambda'(1-x)^3$, the interval of The least upper bound of the set $\left\{\frac{1}{n}, n \in \mathbb{N}\right\}$ is (i) (ii) (iii) (Choose the correct option) (iv) None of the above (Pick up the correct option)

3 .binoo (BCS) STA HC 3/0 2

(b)	Identify	the wrong	statement	

- The intersection of two open sets is open.
- Every open set is an union of open intervals.
- Hone (iii) The union of two open sets is closed.
- (iv) The set of all integers is countable.
- State Bolzano-Weierstrass theorem.
- A sequence cannot converge to more than one limit. (State True or False)
- The value of $\Delta^4(1-x)^4$, the interval of differencing being unity is
 - (i)
 - (ii)
 - _, (iii)
 - (iv) 24

(Choose the correct option)

Lagrange's formula x red (iii) Binomial expansion method (iv) Gauss backward formula (Choose the correct option) (g) If the nth differences of a tabulated function f(x) are constant, the value of independent variables are taken at equal intervals, then f(x) is a polynomial of degree nf(x) is constant theorem f(x) is zero equipment f(x) is a polynomial of degree

(n-1)

3 (Sem-3/CBCS) STA HC 3/G

Given the following data of 1 19 /811A

To interpolate number of workers for

income not exceeding Rs.30 per day,

: 12 32 68 80 100

(Choose the correct option)

DE OH ATE (208 Contd.2) &

Income per day

Workers

not exceeding (Rs.): 10 18

the suitable method is:

oneup (i) Newton's backward formula

3 (Sem -3/CBCS) STA HC 3/G

- Answer the following questions: 2×4=8
 - (a) Using Lagrange's mean value theorem, prove that

$$\left| \frac{1}{\tan^{-1} x} - \tan^{-1} y \right| \le \left| x - y \right| \forall x, y \in \mathbb{R}$$

- Prove that every convergent sequence is bounded. W Lagrange
 - (c) Show that for any real number x,

iv) Gauss be kward for
$$\frac{\pi}{x}$$
 lie option)

- (g) If the n'h differences of a moulated State Taylor's theorem with Lagrange's and Cauchy's form of remainder. qual intervals, then
- 3. Answer any three of the following questions: 5×3=15
 - f(x) is _ instant State and prove Cauchy's first theorem on limits.
 - (b) Expand sinx by Maclaurin's infinite (Choose the correct option)
- 3 (Sem=3/CBCS) STA HC 3/G 4

- State and prove Rolle's theorem.
- (d) If four equidistant values u_{-1} , u_0 , u_1 and u_2 are given and a value u_x is interpolated by Lagrange's formula, show that

$$u_{x} = yu_{0} + xu_{1} + \frac{y(y^{2}-1)}{\Delta^{2}u_{-1}} + \frac{x(x^{2}-1)}{\Delta^{2}u_{0}} \Delta^{2}u_{0}$$
snortesup griwolfo ed 310 (d) 70 (a) 731 and

where x + y = 1 and p and y = 1

- Show that the nth order divided difference of a polynomial of n^{th} degree is constant. who was
- 4. Answer (a) or (b) of the following questions:

(a) (i) Evaluate
$$\lim_{x\to 0} \frac{x-\tan x}{x^3}$$

- (ii) Expand $(1+x)^n$ by Maclaurin's infinite series.
- Prove that a function which is uniformly continuous on an interval is continuous on that interval.

3 (Sem-3/CBCS) STA HC 3/G 5 3 ONE SHEATE BOE Contd.

(ii) Let
$$f(x) =\begin{cases} x^{p} \sin(\frac{1}{x}), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Obtain p such that (i) f(x) is continuous at x = 0 (ii) f(x) is differentiable at x = 0.

Answer (a) or (b) of the following questions:

- State and prove Cauchy's general principle of convergence.
- (b) (i) Solve the difference equation:

$$u_{x+2} - 4u_x = 9x^2 \cos x$$

write a note on use of various interpolation formulae.

Evaluate La : (d) or (b): Answer (a) or (b):

- (a) (i) State Cauchy's nth root test and Leibnitz's test for the convergence of alternating series.
 - (ii) Show that

$$\lim_{n \to \infty} \left[\frac{1}{\sqrt{n^2 + 1}} + \frac{1}{\sqrt{n^2 + 2}} + \dots + \frac{1}{\sqrt{n^2 + n}} \right] = 1$$

3 (Sem -3/CBCS) STA HC 3/G 6 & D\E OH ATS (SOHO\E-Last)

(b) (i) Derive Gauss's interpolation formula for central differences. 5

(iii) State and prove Weddle's rule for numerical integration.

01 001

sile, ratie

3 (Sem-3/CBCS) STA HC 3/G